



Montana Wheat Varieties 2012

Cooperating with the Montana Department of Agriculture
10 W 15th Street, Suite 3100 · Helena, MT 59626

Released: July 16, 2012

WINTER WHEAT

Montana planted 2.2 million acres of winter wheat in 2012, down from 2.25 million acres planted in 2011. Montana is ranked fifth for winter wheat planted acres in the United States, according to the USDA National Agricultural Statistics Service, Montana Field Office. The top four varieties listed below account for over half of the total winter wheat planted in Montana for 2012.


Yellowstone is the most common winter wheat variety for 2012. It accounts for 22.8 percent of the state's planted acreage with 500,500 acres seeded. Yellowstone is a hard red winter wheat developed by the Montana Agricultural Experiment Station and became available in 2005. The variety is high yielding with medium test weight, maturity, height, and protein. It has excellent baking and good noodle quality. Yellowstone is moderately resistant to dwarf smut and stripe rust but is susceptible to stem rust.

Genou is the second most common winter wheat variety planted in Montana. Farmers planted 433,500 acres of the variety, which accounts for 19.7 percent of the total winter wheat acreage in 2012. Genou was released by Montana State University in 2004 and is a solid-stem hard red winter wheat with enhanced yield potential and cold tolerance when compared to Rampart. Test weight, maturity, height, and protein qualities are similar to Rampart and Vanguard.


CDC Falcon is the third leading winter wheat variety planted in Montana in 2012. A total of 154,600 acres were planted accounting for 7.0 percent of the total winter wheat acres planted. CDC Falcon is a semi-dwarf, hard red winter wheat developed by the Crop Development Center in Saskatoon, Saskatchewan and was registered in 1998. It is a high yielding wheat with a short strong straw, good winter-hardiness, and a resistance to stem and leaf rust. This variety has shown average to above average yields on dryland and good performance for irrigated fields.

Rampart is the fourth most common winter variety planted throughout Montana in 2012. A total of 108,500 acres were planted, accounting for 4.9 percent of the state's total acreage. This is a hard red winter wheat that was developed by the Montana Agricultural Experiment Station and released in 1996. It is an awned, red-chaffed, solid-stemmed variety. Kernels are hard red, with a med-sized germ. Rampart is resistant to the wheat stem sawfly and to prevalent races of stem rust. It is susceptible to stripe rust, leaf rust, dwarf smut and the Russian wheat aphid. Rampart has excellent milling and baking qualities.

Montana's Agricultural Districts



2012 Top Seeded Winter Wheat Varieties



Data compiled and summarized by the USDA National Agricultural Statistics Service Montana Field Office as a service provided for the Montana Wheat and Barley Committee.

Steve Anderson, Director
USDA, NASS, MT Field Office

Ron de Yong, Director
Montana Department of Agriculture

Alyssa Peck, Student Intern
USDA, NASS, MT Field Office

SPRING WHEAT

In 2012, Montana producers planted 2.9 million acres of spring wheat, up from 2.45 million acres in 2011. The state remains the second largest spring wheat producer in the country, according to the USDA National Agricultural Statistics Service, Montana Field Office. The top four varieties account for over fifty-five percent of all Montana's 2012 spring wheat acreage.


Vida remains the leading spring wheat variety planted for the second year in a row. Montana producers planted 559,200 acres of Vida this year, accounting for 19.3 percent of the total acres planted. Vida was derived from a cross of Scholar and Reeder in 1998 by the Montana Agricultural Experiment Station and released in 2005. It is a high yielding hard red spring with moderate resistance to leaf and stripe rust. Vida is a semi-dwarf with white glumes and awns, kernels are red ovate with rounded cheeks. Good milling and baking characteristics have made Vida a popular variety each year.

Choteau is the second most common spring wheat variety in 2012 with 447,800 acres accounting for 15.4% of Montana's seeded spring wheat. Choteau was developed by the Montana Agricultural Experiment Station in 2003. This semi-dwarf hard red spring wheat with solid stem prevents sawfly. The spike is lax and tapered with white awns and glumes. Kernels are a red ovate with a medium crease and brush. Choteau is resistant to the prevalent stem and tolerant to the Hessian fly. Yields are similar to McNeal in Montana.

Reeder remains the third most common spring wheat variety planted this year with 389,700 acres planted or 13.4 percent of the seeded spring wheat in Montana. Reeder was developed by the North Dakota Agricultural Experiment Station, released in 1999. It is an awned, semi-dwarf, hard red spring wheat. Reeder is resistant to upper Midwest stem and leaf rust. It was developed by crossing a relative of Stoa and germplasm from Brazil. Its stay-green trait allows for a longer head-filling period and in turn higher yield.

Corbin is the fourth most common Montana spring wheat in 2012 with 202,100 acres seeded accounting for 7.0 percent of the total spring wheat planted. Corbin was developed and released by WestBred LLC, in 2006. It came from a cross between Border and Conan. The line of Corbin is best fitted to the sawfly areas in Montana. The stem and leaves have a slight waxy coating and the chaff is white. It is a semi-dwarf with a resistance to strip rust. Milling and baking quality is acceptable for the market class being grown in Montana.

2012 Top Seeded Spring Wheat Varieties



DURUM WHEAT

Montana producers planted 540,000 acres of durum wheat in 2012, up from 400,000 acres planted in 2011. Montana ranks second for durum acres planted in the United States according to the USDA National Agricultural Statistics Service, Montana Field Office. The top four varieties account for nearly three quarters of the total durum wheat planted in Montana in 2012.


Mountrail has been the leading durum variety planted in the state for the last nine years. This year it was seeded on 143,400 acres or 26.6 percent of Montana's total planted durum acreage. Mountrail is a cross of D8479 and Renville, released by the North Dakota Agricultural Experiment Station in November 1998. This variety is medium height, late maturing and stiff-strawed. It is resistant to both leaf and stem rusts. Mountrail has an above average yield, an average test weight, medium to large kernels, and average protein. It is also known for having a high semolina extract with strong gluten.

Divide is the second most common durum wheat planted in the state this year with 124,900 acres planted—over double that of 2011. It accounts for 23.1 percent of the total durum acres planted in Montana this year. Divide was developed by the North Dakota Agricultural Experiment Station in cooperation with USDA-ARS and released on June 1, 2005. It has high grain yield potential, medium height, strong straw strength, medium maturity and excellent overall quality. Divide is rated as moderately resistant to scab and does provide slightly enhanced protection against scab relative to other durum varieties.

Strongfield became the third most common durum variety planted in 2012. There were 85,600 acres seeded, accounting for 15.9 percent of Montana's planted acreage. Strongfield was released by Ag Canada in 2004. It is medium in height and maturity. Strongfield has good resistance to leaf and stem rusts but only moderately resistant to common bunt. The variety exhibits good end use qualities by having high protein, low grain cadmium concentration, high yellow pigment, and a moderately strong gluten content.

Alzada is ranked as the fourth most common durum variety with 41,200 acres planted in 2012, or 7.6 percent of Montana's total acres. Alzada was developed by WestBred, LLC in Bozeman, Montana and first released for commercial planting in 2004. The variety has competitive yields along with excellent straw strength and good sawfly tolerance. Alzada has a high test weight, protein content, and demonstrates good scores for semolina color and gluten strength.

2012 Top Seeded Durum Wheat Varieties



Winter Wheat: 2012 Seeded Acreage and Percent of Total Seeded by Districts

Variety	Northwest		North Central		Northeast		Central		Southwest		South Central		Southeast		State Total	
	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%
Yellowstone	2.6	17.4	222.0	18.3	46.3	22.8	138.2	32.9	10.5	38.9	60.8	30.4	20.1	16.5	500.5	22.8
Genou	--	--	378.5	31.2	1.8	0.9	39.1	9.3	2.7	10.1	10.4	5.2	1.0	0.8	433.5	19.7
CDC Falcon	--	--	104.3	8.6	5.5	2.7	41.6	9.9	1.2	4.3	1.6	0.8	0.4	0.3	154.6	7.0
Rampart	--	--	104.3	8.6	--	--	3.8	0.9	--	--	--	--	0.4	0.3	108.5	4.9
Ledger	--	--	88.5	7.3	--	--	14.3	3.4	--	--	--	--	5.0	4.1	107.8	4.9
Jerry	--	--	2.4	0.2	61.3	30.2	8.8	2.1	--	--	0.2	0.1	35.0	28.7	107.7	4.9
AP503 CL2	0.4	2.9	20.6	1.7	1.4	0.7	43.3	10.3	2.0	7.3	29.6	14.8	1.0	0.8	98.3	4.5
Hawken	--	--	26.7	2.2	4.9	2.4	17.6	4.2	--	--	3.8	1.9	0.4	0.3	53.4	2.4
Norris	--	--	34.0	2.8	--	--	5.5	1.3	0.4	1.5	6.8	3.4	0.4	0.3	47.1	2.1
Pryor	--	--	23.0	1.9	--	--	13.9	3.3	0.8	2.9	4.6	2.3	1.5	1.2	43.8	2.0
Jagalene	0.5	3.6	7.3	0.6	2.6	1.3	18.5	4.4	--	--	8.0	4.0	3.9	3.2	40.8	1.9
Carter	--	--	40.0	3.3	--	--	--	--	--	--	--	--	--	--	40.0	1.8
Morgan	--	--	30.3	2.5	7.5	3.7	--	--	--	--	--	--	--	--	37.8	1.7
Decade	0.1	0.5	7.3	0.6	19.5	9.6	4.2	1.0	--	--	0.8	0.4	2.1	1.7	34.0	1.5
WB Quake	--	--	19.4	1.6	--	--	5.5	1.3	3.2	11.7	2.4	1.2	--	--	30.5	1.4
Tiber	--	--	--	--	3.0	1.5	3.4	0.8	0.7	2.5	8.6	4.3	6.8	5.6	22.5	1.0
Willow Creek	0.1	0.8	1.2	0.1	4.5	2.2	7.1	1.7	--	--	3.0	1.5	3.1	2.5	19.0	0.9
Radiant	--	--	14.6	1.2	3.0	1.5	--	--	--	--	--	--	1.1	0.9	18.7	0.9
Neeley	--	--	2.4	0.2	0.8	0.4	14.7	3.5	0.4	1.4	0.2	0.1	--	--	18.5	0.8
Vanguard	--	--	8.5	0.7	--	--	5.0	1.2	--	--	2.8	1.4	--	--	16.3	0.7
Promontory	0.6	3.9	--	--	--	--	1.7	0.4	--	--	8.6	4.3	0.5	0.4	11.4	0.5
Overland	--	--	--	--	4.7	2.3	--	--	--	--	--	--	5.9	4.8	10.6	0.5
Redwin	--	--	2.4	0.2	--	--	--	--	0.4	1.4	--	--	5.7	4.7	8.5	0.4
Other & Unknown	10.7	70.9	75.3	6.2	36.2	17.8	33.8	8.1	4.7	18.0	47.8	23.9	27.7	22.9	236.2	10.8
All Varieties	15.0	100.0	1,213.0	100.0	203.0	100.0	420.0	100.0	27.0	100.0	200.0	100.0	122.0	100.0	2,200.0	100.0

--Data not published for given year.

Winter Wheat: 2011 Seeded Acreage and Percent of Total Seeded by Districts ^{1/}

Variety	Northwest		North Central		Northeast		Central		Southwest		South Central		Southeast		State Total	
	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%
Genou	--	--	560.7	43.1	2.4	1.6	44.8	10.4	3.7	14.7	15.4	7.1	1.3	1.2	628.3	27.9
Yellowstone	0.4	3.1	147.0	11.3	64.0	42.2	75.4	17.5	8.0	32.1	45.7	21.1	40.4	36.1	380.9	16.9
CDC Falcon	0.4	3.2	119.7	9.2	4.4	2.9	46.5	10.8	2.7	10.9	6.5	3.0	0.2	0.2	180.4	8.0
Rampart	--	--	114.5	8.8	--	--	4.3	1.0	--	--	2.4	1.1	--	--	121.2	5.4
Ledger	--	--	89.8	6.9	--	--	24.6	5.7	--	--	1.9	0.9	--	--	116.3	5.2
AP503 CL2	0.5	3.8	14.3	1.1	0.2	0.1	34.9	8.1	1.5	5.8	59.8	27.6	2.1	1.9	113.3	5.0
Jerry	0.5	3.6	6.5	0.5	36.0	23.7	15.5	3.6	--	--	--	--	11.7	10.5	70.2	3.1
Jagalene	0.9	6.8	9.1	0.7	--	--	37.9	8.8	--	--	12.1	5.6	7.4	6.6	67.4	3.0
Pryor	--	--	23.4	1.8	--	--	27.6	6.4	1.8	7.2	5.6	2.6	6.0	5.4	64.4	2.9
Hawken	--	--	9.1	0.7	2.1	1.4	38.4	8.9	--	--	--	--	0.8	0.7	50.4	2.3
Norris	0.2	1.9	36.4	2.8	--	--	2.2	0.5	1.0	3.8	9.5	4.4	--	--	49.3	2.2
Morgan	--	--	36.4	2.8	11.2	7.4	--	--	--	--	0.4	0.2	--	--	48.0	2.1
Neeley	--	--	7.8	0.6	0.3	0.2	18.1	4.2	1.3	5.0	8.9	4.1	2.1	1.9	38.5	1.7
Vanguard	--	--	15.6	1.2	--	--	14.7	3.4	--	--	--	--	--	--	30.3	1.4
Carter	--	--	26.0	2.0	--	--	0.4	0.1	--	--	--	--	--	--	26.4	1.2
Tiber	--	--	3.9	0.3	2.1	1.4	--	--	0.5	1.8	9.3	4.3	5.1	4.6	20.9	0.9
Willow Creek	0.4	3.2	2.6	0.2	5.0	3.3	6.0	1.4	--	--	0.9	0.4	0.8	0.7	15.7	0.7
Bynum	--	--	9.1	0.7	--	--	0.4	0.1	--	--	1.1	0.5	--	--	10.6	0.5
Redwin	--	--	--	--	2.4	1.6	2.6	0.6	--	--	2.6	1.2	1.9	1.7	9.5	0.4
Promontory	1.5	11.6	--	--	--	--	2.2	0.5	--	--	5.6	2.6	--	--	9.3	0.4
Rocky	--	--	9.1	0.7	--	--	--	--	--	--	--	--	--	--	9.1	0.4
Winalta	--	--	2.6	0.2	2.0	1.3	--	--	--	--	--	--	2.7	2.4	7.3	0.3
Carlisle	0.1	0.8	5.2	0.4	--	--	--	--	--	--	--	--	--	--	5.3	0.2
Other & Unknown	8.1	62.0	52.2	4.0	19.6	12.9	34.5	8.0	4.5	18.7	28.8	13.3	29.3	26.1	177.0	7.9
All Varieties	13.0	100.0	1,301.0	100.0	151.7	100.0	431.0	100.0	25.0	100.0	216.5	100.0	111.8	100.0	2,250.0	100.0

^{1/} Revised. --Data not published for given year.

Spring Wheat: 2012 Seeded Acreage and Percent of Total Seeded by Districts

Variety	Northwest		North Central		Northeast		Central		Southwest		South Central		Southeast		State Total	
	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%
Vida	0.2	0.5	183.3	17.0	332.0	25.0	26.9	13.6	1.1	1.9	4.1	6.1	11.6	8.5	559.2	19.3
Choteau	--	--	247.9	23.0	155.4	11.7	30.3	15.3	3.6	6.0	4.8	7.1	5.8	4.3	447.8	15.4
Reeder	--	--	9.7	0.9	340.0	25.6	9.3	4.7	--	--	5.0	7.4	25.7	18.9	389.7	13.4
Corbin	--	--	198.4	18.4	2.7	0.2	1.0	0.5	--	--	--	--	--	--	202.1	7.0
ONeal	--	--	120.7	11.2	27.9	2.1	39.2	19.8	0.5	0.9	1.6	2.3	--	--	189.9	6.5
McNeal	--	--	37.7	3.5	89.0	6.7	11.1	5.6	--	--	4.6	6.7	14.3	10.5	156.7	5.4
Conan	--	--	66.8	6.2	1.3	0.1	0.4	0.2	--	--	--	--	0.7	0.5	69.2	2.4
Brennan	0.8	2.5	1.1	0.1	43.8	3.3	5.7	2.9	1.7	2.8	0.3	0.4	1.5	1.1	54.9	1.9
WB Gunnison	--	--	48.5	4.5	--	--	4.4	2.2	--	--	--	--	0.3	0.2	53.2	1.8
Mott	--	--	5.4	0.5	37.2	2.8	--	--	--	--	--	--	7.1	5.2	49.7	1.7
Fortuna	--	--	29.1	2.7	--	--	14.9	7.5	0.2	0.3	4.6	6.8	--	--	48.8	1.7
Kelby	1.4	4.3	--	--	35.9	2.7	6.9	3.5	0.3	0.5	1.4	2.0	2.2	1.6	48.1	1.7
Glenn	--	--	--	--	29.2	2.2	3.0	1.5	--	--	--	--	3.3	2.4	35.5	1.2
Solano	13.2	41.3	--	--	2.7	0.2	5.1	2.6	5.7	9.5	6.7	9.8	1.2	0.9	34.6	1.2
AC Lillian	--	--	5.4	0.5	27.9	2.1	--	--	--	--	1.1	1.6	--	--	34.4	1.2
SY Tyra	--	--	21.6	2.0	1.3	0.1	3.2	1.6	--	--	2.7	4.0	1.6	1.2	30.4	1.0
Amidon	--	--	2.2	0.2	23.9	1.8	--	--	--	--	0.1	0.2	--	--	26.2	0.9
Hank	2.7	8.4	1.1	0.1	1.3	0.1	10.9	5.5	1.7	2.9	4.5	6.6	2.7	2.0	24.9	0.9
Steele	0.2	0.5	1.1	0.1	8.0	0.6	0.4	0.2	--	--	--	--	14.1	10.4	23.8	0.8
Westbred 936	--	--	9.7	0.9	1.3	0.1	4.2	2.1	3.1	5.2	5.1	7.5	--	--	23.4	0.8
Ernest	--	--	17.2	1.6	2.7	0.2	0.2	0.1	--	--	--	--	1.5	1.1	21.6	0.7
Volt	0.4	1.1	6.5	0.6	8.0	0.6	3.0	1.5	0.2	0.4	1.3	1.9	0.3	0.2	19.7	0.7
AP 605 CL	0.1	0.2	5.4	0.5	9.3	0.7	1.6	0.8	0.7	1.1	1.6	2.4	--	--	18.7	0.6
Jedd	--	--	9.7	0.9	1.3	0.1	2.4	1.2	0.5	0.9	0.1	0.2	0.3	0.2	14.3	0.5
Other & Unknown	13.0	41.2	49.5	4.6	145.9	11.0	13.9	7.1	40.7	67.6	18.4	27.0	41.8	30.8	323.2	11.3
All Varieties	32.0	100.0	1,078.0	100.0	1,328.0	100.0	198.0	100.0	60.0	100.0	68.0	100.0	136.0	100.0	2,900.0	100.0

--Data not published for given year.

Spring Wheat: 2011 Seeded Acreage and Percent of Total Seeded by Districts ^{1/}

Variety	Northwest		North Central		Northeast		Central		Southwest		South Central		Southeast		State Total	
	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%	(000)	%
Vida	0.4	1.1	191.6	19.9	322.8	29.7	26.4	16.5	0.8	1.3	5.1	7.7	6.6	8.1	553.7	22.6
Choteau	0.1	0.3	253.3	26.3	142.4	13.1	18.9	11.8	7.1	12.0	8.3	12.5	13.2	16.1	443.3	18.1
Reeder	0.3	0.9	1.9	0.2	252.2	23.2	11.8	7.4	--	--	4.7	7.1	8.7	10.6	279.6	11.4
Corbin	--	--	175.3	18.2	4.3	0.4	1.4	0.9	--	--	0.7	1.1	--	--	181.7	7.4
McNeal	--	--	23.1	2.4	100.0	9.2	12.2	7.6	1.7	2.9	2.6	4.0	2.5	3.1	142.1	5.8
ONeal	0.2	0.7	89.6	9.3	13.0	1.2	28.0	17.5	0.7	1.2	1.5	2.3	1.8	2.2	134.8	5.5
Fortuna	--	--	57.8	6.0	--	--	17.8	11.1	--	--	4.2	6.3	--	--	79.8	3.3
Conan	--	--	67.4	7.0	6.5	0.6	--	--	--	--	0.3	0.5	--	--	74.2	3.0
AC Lillian	--	--	9.6	1.0	30.4	2.8	--	--	--	--	1.0	1.5	--	--	41.0	1.7
Kelby	3.6	10.8	--	--	18.5	1.7	4.2	2.6	0.6	1.1	1.1	1.7	3.4	4.1	31.4	1.3
Amidon	--	--	--	--	25.0	2.3	--	--	--	--	--	--	0.2	0.3	25.2	1.0
Hank	2.3	7.0	--	--	6.5	0.6	9.1	5.7	3.1	5.2	2.7	4.1	--	--	23.7	1.0
Steele	--	--	--	--	4.3	0.4	0.2	0.1	--	--	--	--	18.1	22.1	22.6	0.9
Solano	9.8	29.7	--	--	--	--	0.2	0.1	2.0	3.4	6.8	10.3	2.2	2.7	21.0	0.9
Kuntz	2.0	6.0	2.9	0.3	6.5	0.6	5.1	3.2	3.6	6.1	--	--	--	--	20.1	0.8
Ernest	--	--	15.4	1.6	4.3	0.4	--	--	--	--	--	--	--	--	19.7	0.8
Westbred 936	0.2	0.5	3.9	0.4	--	--	4.5	2.8	9.1	15.5	0.5	0.7	--	--	18.2	0.7
WB Gunnison	--	--	17.3	1.8	--	--	--	--	--	--	--	--	--	--	17.3	0.7
Glenn	--	--	--	--	15.2	1.4	--	--	1.4	2.3	--	--	--	--	16.6	0.7
Brennan	0.3	0.8	--	--	12.0	1.1	0.2	0.1	0.8	1.3	0.9	1.4	--	--	14.2	0.6
Mott	1.0	3.0	2.9	0.3	9.8	0.9	--	--	--	--	--	--	--	--	13.7	0.6
Jedd	0.2	0.6	5.8	0.6	--	--	3.5	2.2	--	--	4.0	6.0	--	--	13.5	0.6
Freyr	--	--	1.9	0.2	6.5	0.6	--	--	1.4	2.4	3.0	4.6	0.5	0.6	13.3	0.5
Scholar	--	--	--	--	10.9	1.0	0.5	0.3	--	--	--	--	0.2	0.3	11.6	0.5
Other & Unknown	12.6	38.6	43.3	4.5	95.9	8.8	16.0	10.1	26.7	45.3	18.6	28.2	24.6	29.8	237.7	9.6
All Varieties	33.0	100.0	963.0	100.0	1,087.0	100.0	160.0	100.0	59.0	100.0	66.0	100.0	82.0	100.0	2,450.0	100.0

^{1/} Revised. --Data not published for given year.

Winter Wheat: Percent of Total Seeded Acreage 2006-2012 & Seeded Acreage 2011-2012

Variety	Percent of Total Seeded Acreage							Seeded Acres (000)	
	2006	2007	2008	2009	2010	2011 1/	2012	2011 1/	2012
Yellowstone	--	3.7	8.2	12.8	18.1	16.9	22.8	380.9	500.5
Genou	2.2	15.8	19.5	24.4	30.2	27.9	19.7	628.3	433.5
CDC Falcon	10.9	13.3	13.2	9.0	7.4	8.0	7.0	180.4	154.6
Rampart	19.2	9.6	7.4	5.6	7.4	5.4	4.9	121.2	108.5
Ledger	2.4	5.8	6.8	6.5	4.2	5.2	4.9	116.3	107.8
Jerry	--	0.7	2.9	3.6	1.7	3.1	4.9	70.2	107.7
AP503 CL2	--	--	--	--	1.2	5.0	4.5	113.3	98.3
Hawken	--	--	--	--	--	2.3	2.4	50.4	53.4
Norris	--	--	0.5	0.9	2.5	2.2	2.1	49.3	47.1
Pryor	3.3	4.6	3.5	2.3	2.9	2.9	2.0	64.4	43.8
Jagalene	6.2	5.6	5.7	5.5	5.0	3.0	1.9	67.4	40.8
Carter	--	--	--	0.5	1.2	1.2	1.8	26.4	40.0
Morgan	7.1	5.3	4.6	4.6	2.5	2.1	1.7	48.0	37.8
Decade	--	--	--	--	--	--	1.5	--	34.0
WB Quake	--	--	--	--	--	--	1.4	--	30.5
Tiber	7.1	5.0	5.6	4.0	2.2	0.9	1.0	20.9	22.5
Willow Creek	--	--	--	0.4	0.7	0.7	0.9	15.7	19.0
Radiant	--	--	--	--	0.4	--	0.9	--	18.7
Neeley	9.3	7.8	4.2	2.2	2.4	1.7	0.8	38.5	18.5
Vanguard	2.9	1.4	1.7	0.8	1.1	1.4	0.7	30.3	16.3
Promontory	3.5	2.5	0.8	2.1	1.7	0.4	0.5	9.3	11.4
Overland	--	--	--	--	--	--	0.5	--	10.6
Redwin	1.7	1.4	1.0	1.5	0.5	0.4	0.4	9.5	8.5
Bynum	--	--	--	0.5	0.4	0.5	--	10.6	--
Carlisle	--	--	0.6	0.6	0.3	0.2	--	5.3	--
Rocky	7.1	4.4	2.1	0.7	1.0	0.4	--	9.1	--
Winalta	1.0	0.8	0.4	0.4	0.3	0.3	--	7.3	--
Other & Unknown	16.1	12.3	11.3	11.1	4.7	7.9	10.8	177.0	236.2
All Varieties	100.0	100.0	100.0	100.0	100.0	100.0	100.0	2,250.0	2,200.0

1/ Revised. --Data not published for given year.

Spring Wheat: Percent of Total Seeded Acreage 2006-2012 & Seeded Acreage 2011-2012

Variety	Percent of Total Seeded Acreage							Seeded Acres (000)	
	2006	2007	2008	2009	2010	2011 1/	2012	2011 1/	2012
Vida	--	--	--	5.0	12.8	22.6	19.3	553.7	559.2
Choteau	16.5	22.4	21.6	22.7	23.0	18.1	15.4	443.3	447.8
Reeder	27.6	28.1	25.0	24.0	16.5	11.4	13.4	279.6	389.7
Corbin	--	2.2	3.4	4.0	5.6	7.4	7.0	181.7	202.1
ONeal	--	--	--	1.4	4.2	5.5	6.5	134.8	189.9
McNeal	21.3	14.9	12.1	10.6	8.4	5.8	5.4	142.1	156.7
Conan	6.8	5.0	2.9	3.2	2.9	3.0	2.4	74.2	69.2
Brennan	--	--	--	--	--	0.6	1.9	14.2	54.9
WB Gunnison	--	--	--	--	--	0.7	1.8	17.3	53.2
Mott	--	--	--	--	--	0.6	1.7	13.7	49.7
Fortuna	4.1	4.3	4.1	2.5	3.8	3.3	1.7	79.8	48.8
Kelby	--	--	1.0	1.0	1.1	1.3	1.7	31.4	48.1
Glenn	--	0.6	--	--	0.6	0.7	1.2	16.6	35.5
Solano	--	--	--	0.8	0.6	0.9	1.2	21.0	34.6
AC Lillian	--	--	2.9	1.6	2.5	1.7	1.2	41.0	34.4
SY Tyra	--	--	--	--	--	--	1.0	--	30.4
Amidon	1.8	1.7	1.7	1.0	0.5	1.0	0.9	25.2	26.2
Hank	2.3	1.9	2.6	1.7	1.5	1.0	0.9	23.7	24.9
Steele	--	--	--	0.5	0.4	0.9	0.8	22.6	23.8
Westbred 936	0.7	0.9	0.8	1.3	0.7	0.7	0.8	18.2	23.4
Ernest	5.5	3.1	2.3	1.5	1.6	0.8	0.7	19.7	21.6
Volt	--	--	--	--	--	--	0.7	--	19.7
AP 605 CL	--	--	--	--	--	--	0.6	--	18.7
Jedd	--	--	--	0.8	0.8	0.6	0.5	13.5	14.3
Kuntz	--	--	--	1.1	0.8	0.8	--	20.1	--
Freyr	--	0.7	0.6	1.0	0.9	0.5	--	13.3	--
Scholar	0.4	1.2	0.7	1.2	0.4	0.5	--	11.6	--
Other & Unknown	13.0	13.0	18.3	13.1	10.4	9.6	11.3	237.7	323.2
All Varieties	100.0	100.0	100.0	100.0	100.0	100.0	100.0	2,450.0	2,900.0

1/ Revised. --Data not published for given year.

Durum Wheat: 2012 Seeded Acreage and Percent of Total Seeded by Districts

Variety	North Central		Northeast		Other Districts		State Total	
	(000)	%	(000)	%	(000)	%	(000)	%
Mountrail	--	--	142.1	29.6	1.3	11.2	143.4	26.6
Divide	--	--	120.5	25.1	4.4	36.4	124.9	23.1
Strongfield	7.8	16.2	77.8	16.2	--	--	85.6	15.9
Alzada	29.6	61.6	11.0	2.3	0.6	5.0	41.2	7.6
Alkabo	1.1	2.2	19.2	4.0	1.4	11.7	21.7	4.0
Kyle	--	--	17.3	3.6	--	--	17.3	3.2
Grenora	--	--	15.8	3.3	--	--	15.8	2.9
Lebsock	--	--	12.5	2.6	--	--	12.5	2.3
Pierce	--	--	7.7	1.6	--	--	7.7	1.4
Westhope	--	--	7.2	1.5	--	--	7.2	1.3
Dilse	--	--	6.7	1.4	--	--	6.7	1.2
Tioga	1.7	3.5	4.8	1.0	--	--	6.5	1.2
Vic	--	--	4.8	1.0	--	--	4.8	0.9
AC Avonlea	1.0	2.1	1.4	0.3	--	--	2.4	0.4
Medora	--	--	2.4	0.5	--	--	2.4	0.4
Other & Unknown	6.8	14.4	28.8	6.0	4.3	35.7	39.9	7.6
All Varieties	48.0	100.0	480.0	100.0	12.0	100.0	540.0	100.0

--Data not published for given year.

Durum Wheat: 2011 Seeded Acreage and Percent of Total Seeded by Districts ^{1/}

Variety	North Central		Northeast		Other Districts		State Total	
	(000)	%	(000)	%	(000)	%	(000)	%
Mountrail	--	--	110.5	31.3	--	--	110.5	27.6
Strongfield	5.3	13.5	53.7	15.2	0.6	7.5	59.6	14.9
Divide	1.0	2.6	47.3	13.4	--	--	48.3	12.1
Alzada	28.0	71.9	8.8	2.5	0.5	5.7	37.3	9.3
Lebsock	--	--	20.1	5.7	--	--	20.1	5.0
Kyle	0.8	2.0	19.1	5.4	--	--	19.9	5.0
Alkabo	--	--	18.7	5.3	--	--	18.7	4.7
Grenora	--	--	16.2	4.6	--	--	16.2	4.1
Westhope	--	--	8.5	2.4	--	--	8.5	2.1
Pierce	--	--	7.8	2.2	--	--	7.8	2.0
Vic	--	--	4.9	1.4	--	--	4.9	1.2
Dilse	--	--	1.8	0.5	2.9	35.8	4.7	1.2
Ward	--	--	3.2	0.9	--	--	3.2	0.8
AC Morse	2.1	5.3	--	--	--	--	2.1	0.5
Monroe	--	--	1.4	0.4	--	--	1.4	0.4
Other & Unknown	1.8	4.7	31.0	8.8	4.0	51.0	36.8	9.1
All Varieties	39.0	100.0	353.0	100.0	8.0	100.0	400.0	100.0

^{1/} Revised. -- Data not published for given year.

Durum Wheat: Percent Total Seeded Acreage 2006-2012 & Seeded Acreage 2011-2012

Variety	Percent of Total Seeded Acreage						Seeded Acres (000)		
	2006	2007	2008	2009	2010	2011 ^{1/}	2012	2011 ^{1/}	2012
Mountrail	52.6	44.8	40.8	34.6	25.2	27.6	26.6	110.5	143.4
Divide	--	0.7	4.0	5.5	10.1	12.1	23.1	48.3	124.9
Strongfield	--	1.2	8.6	7.4	13.4	14.9	15.9	59.6	85.6
Alzada	4.8	5.1	6.4	7.3	9.2	9.3	7.6	37.3	41.2
Alkabo	--	--	1.5	4.6	4.6	4.7	4.0	18.7	21.7
Kyle	15.7	15.4	14.2	8.7	8.3	5.0	3.2	19.9	17.3
Grenora	--	0.6	4.0	6.5	5.9	4.1	2.9	16.2	15.8
Lebsock	10.1	12.3	6.0	7.7	6.0	5.0	2.3	20.1	12.5
Pierce	5.6	5.3	2.7	1.9	1.2	2.0	1.4	7.8	7.7
Westhope	--	--	--	--	--	2.1	1.3	8.5	7.2
Dilse	0.8	2.8	0.8	0.6	3.0	1.2	1.2	4.7	6.7
Tioga	--	--	--	--	--	--	1.2	--	6.5
Vic	0.9	0.4	--	--	0.5	1.2	0.9	4.9	4.8
AC Avonlea	2.3	1.0	1.3	3.1	1.8	--	0.4	--	2.4
Medora	--	--	--	--	--	--	0.4	--	2.4
Monroe	0.6	1.3	2.1	2.3	0.9	0.4	--	1.4	--
Ward	2.3	1.5	--	0.6	--	0.8	--	3.2	--
AC Morse	--	--	0.3	0.9	--	0.5	--	2.1	--
Other & Unknown	4.3	7.6	7.3	8.3	9.9	9.1	7.6	36.8	39.9
All Varieties	100.0	100.0	100.0	100.0	100.0	100.0	100.0	400.0	540.0

^{1/} Revised. --Data not published for given year.